

Irreducible pseudo 2-factor isomorphic cubic bipartite graphs

M. Abreu,

Dipartimento di Matematica, Università della Basilicata,

C. da Macchia Romana, 85100 Potenza, Italy.

e-mail: marien.abreu@unibas.it

D. Labbate,

Dipartimento di Matematica, Politecnico di Bari

I-70125 Bari, Italy.

e-mail: labbate@poliba.it

J. Sheehan,

Department of Mathematical Sciences, King's College,

Old Aberdeen AB24 3UE, Scotland.

e-mail: j.sheehan@maths.abdn.ac.uk

Abstract

A bipartite graph is *pseudo 2-factor isomorphic* if all its 2-factors have the same parity of number of circuits. In [2] we proved that the only essentially 4-edge-connected pseudo 2-factor isomorphic cubic bipartite graph of girth 4 is $K_{3,3}$, and conjectured [2, Conjecture 3.6] that the only essentially 4-edge-connected cubic bipartite graphs are $K_{3,3}$, the Heawood graph and the Pappus graph.

There exists a characterization of symmetric configurations n_3 due to Martinetti (1886) in which all symmetric configurations n_3 can be obtained from an infinite set of so called *irreducible* configurations [10]. The list of irreducible configurations has been completed by Boben [4] in terms of their *irreducible Levi graphs*.

In this paper we characterize irreducible pseudo 2-factor isomorphic cubic bipartite graphs proving that the only pseudo 2-factor isomorphic irreducible Levi graphs are the Heawood and Pappus graphs. Moreover, the obtained characterization allows us to partially prove the above Conjecture.

1 Introduction

All graphs considered are finite and simple (without loops or multiple edges). A graph with a 2-factor is said to be *2-factor hamiltonian* if all its 2-factors are Hamilton circuits, and, more generally, *2-factor isomorphic* if all its 2-factors are isomorphic. Examples of such graphs are K_4 , K_5 , $K_{3,3}$, the Heawood graph (which are all 2-factor hamiltonian) and the Petersen graph (which is 2-factor isomorphic). Several recent papers have addressed the problem of characterizing families of graphs (particularly regular graphs) which have these properties. It is shown in [3, 7] that k -regular 2-factor isomorphic bipartite graphs exist only when $k \in \{2, 3\}$ and an infinite family of 3-regular 2-factor hamiltonian bipartite graphs, based on $K_{3,3}$ and the Heawood graph, is constructed in [7]. It is conjectured in [7] that every 3-regular 2-factor hamiltonian bipartite graph belongs to this family. Faudree, Gould and Jacobsen in [6] determine the maximum number of edges in both 2-factor hamiltonian graphs and 2-factor hamiltonian bipartite graphs. In addition, Diwan [5] has shown that K_4 is the only 3-regular 2-factor hamiltonian planar graph.

Moreover, 2-factor isomorphic bipartite graphs are extended in [2] to the more general family of *pseudo 2-factor isomorphic graphs* i.e. graphs G with the property that the parity of the number of circuits in a 2-factor is the same for all 2-factors of G . Example of these graphs are $K_{3,3}$, the Heawood graph H_0 and the Pappus graph P_0 . Finally, it is proved in [2] that pseudo 2-factor isomorphic $2k$ -regular graphs and k -regular digraphs do not exist for $k \geq 4$. Recently these results has been generalized in [1] for regular graphs which are not necessarily bipartite.

An incidence structure is *linear* if two different points are incident with at most one line. A *symmetric configuration* n_k (or n_k *configuration*) is a linear incidence structure consisting of n points and n lines such that each point and line is respectively incident with k lines and points. Let $\mathcal{C}$ be a symmetric configuration n_k , its *Levi graph* $G(\mathcal{C})$ is a k -regular bipartite graph whose vertex set are the points and the lines of $\mathcal{C}$ and there is an edge between a point and a line in the graph if and only if they are incident in $\mathcal{C}$. We will indistinctly refer to Levi graphs of configurations as their *incidence graphs*.

Let G be a graph and E_1 be an edge-cut of G . We say that E_1 is a *non-trivial edge-cut* if all components of $G - E_1$ have at least two vertices. The graph G is *essentially 4-edge-connected* if G is 3-edge-connected and has no non-trivial 3-edge-cuts.

Conjecture 1.1 [2] *Let G be an essentially 4-edge-connected pseudo 2-factor isomorphic cubic bipartite graph. Then $G \in \{K_{3,3}, H_0, P_0\}$.*

Theorem 1.2 [2] *Let G be an essentially 4-edge-connected pseudo 2-factor isomorphic cubic bipartite graph. Suppose that G contains a 4-circuit. Then $G = K_{3,3}$.* $\square$

It follows from Theorem 1.2 that an essentially 4-edge-connected pseudo 2-factor isomorphic cubic bipartite graph of girth greater than or equal to 6 is the Levi graph of a symmetric configuration n_3 . In 1886 V. Martinetti [10] characterized symmetric configurations n_3 , showing that they can be obtained from an infinite set of so called *irreducible* configurations, of which he gave a list. Recently, Boben proved that Martinetti's list of irreducible configurations was incomplete and completed it [4]. Boben's list of irreducible configurations was obtained characterizing their Levi graphs, which he called *irreducible Levi graphs* (cf. Section 2).

In this paper, we characterize *irreducible* pseudo 2-factor isomorphic cubic bipartite graphs proving that *the Heawood and the Pappus graphs are the only irreducible Levi graphs which are pseudo 2-factor isomorphic cubic bipartite*. Moreover, the obtained characterization allows us to partially prove Conjecture 1.1, i.e. in the case of irreducible pseudo 2-factor isomorphic cubic bipartite graphs.

2 Symmetric Configurations n_3

In 1886, Martinetti [10] provided a construction for a symmetric configuration n_3 from a symmetric configuration $(n-1)_3$, say $\mathcal{C}$. Suppose that in $\mathcal{C}$ there exist two parallel (non-intersecting) lines $l_1 = \{\alpha, \alpha_1, \alpha_2\}$ and $r_1 = \{\beta, \beta_1, \beta_2\}$ such that the points α and β are not on a common line. Then a symmetric configuration n_3 is obtained from $\mathcal{C}$ by deleting the lines l_1, r_1 , adding a point μ and adding the lines $h_1 = \{\mu, \alpha_1, \alpha_2\}$, $h_2 = \{\mu, \beta_1, \beta_2\}$ and $h_3 = \{\mu, \alpha, \beta\}$. Not all symmetric configurations n_3 can be obtained using this method on some symmetric configuration $(n-1)_3$. The configurations that cannot be obtained in this way are called *irreducible configurations*, while the others are *reducible configurations*. However, if all irreducible symmetric configurations n_3 are known, then all symmetric configurations n_3 can be constructed iteratively with Martinetti's method. The list of irreducible configurations in [10] turned out to be incomplete and it has been recently completed by Boben in [4, Thm. 8].

Theorem 2.1 [4] *All connected irreducible n_3 configurations are:*

1. *cyclic configurations with base line $\{0, 1, 3\}$;*
2. *the configurations with their incidence graphs $T_1(n)$, $T_2(n)$, $T_3(n)$, $n \geq 1$, each of them giving precisely one $(10n)_3$ configuration, and;*

3. the Pappus configuration.

As mentioned before, Boben's list was obtained by studying the Levi graphs of irreducible configurations, which are called *irreducible Levi graphs*. Such graphs turned out to be either the Pappus graph, or belong to one of four infinite families $D(n)$, $T_1(n)$, $T_2(n)$, $T_3(n)$, $n \geq 1$, which we now proceed to describe.

The $D(n)$ family: Let $C(n)$, $n \geq 1$, be the graph on $6n$ vertices, consisting of n segments (6-circuits labeled as in Fig. 1), linked by the edges $v_{i-1}^1 u_i^1$, $v_{i-1}^2 u_i^4$, and $u_{i-1}^3 u_i^2$, for $i \geq 2$.




Figure 1: label for the 6-circuits of the definition of $C(n)$

Let the graph $D(n)$, $n \geq 1$ be defined as follows:

For $n \equiv 0 \pmod{3}$ let $D(n)$ be the graph $C(m)$, $m = n/3$, with the edges $u_1^1 v_m^1$, $u_1^4 v_m^2$, and $u_1^2 u_m^3$ added (cf. Fig. 2).

For $n \equiv 1 \pmod{3}$ let $D(n)$ be the graph $C(m)$, $m = (n-1)/3$, with two vertices w_m^1, w_m^2 and the edges $u_1^1 w_m^1$, $u_1^2 v_m^2$, $u_1^4 w_m^2$, $w_m^1 w_m^2$, $w_m^1 u_m^3$, $w_m^2 v_m^1$ added.

For $n \equiv 2 \pmod{3}$ let $D(n)$ be the graph $C(m)$, $m = (n-2)/3$, with four vertices $w_m^1, w_m^2, w_m^3, w_m^4$ and the edges $v_m^1 w_m^1$, $v_m^2 w_m^4$, $u_m^3 w_m^2$, $u_1^1 w_m^4$, $u_1^2 w_m^1$, $u_1^4 w_m^3$, $w_m^1 w_m^2$, $w_m^2 w_m^3$, $w_m^3 w_m^4$ added.




Figure 2: The graph $D(9)$

A *cyclic configuration* has $\mathbb{Z}_n = \{0, 1, \dots, n-1\}$ as set of points and $\mathcal{B} = \{\{0, b, c\}, \{1, b+1, c+1\}, \dots, \{n-1, b+n-1, c+n-1\}\}$ as set of lines, where the operations are modulo n , and the *base line* is $\{0, b, c\}$ for $b, c \in \mathbb{Z}_n$.

Note that the graphs $D(n)$ are the Levi graphs of the cyclic n_3 configurations with base line $\{0, 1, 3\}$. In particular, for $n = 7$ the cyclic 7_3 configuration is the Fano plane and $D(7)$ is the Heawood graph H_0 .

The $T_1(n)$, $T_2(n)$ and $T_3(n)$ families: Let $T(n)$, $n \geq 1$, be the graph on $20n$ vertices consisting of n segments G_T shown in Fig. 3, linked by the edges $v_{i-1}^1 u_i^1$, $v_{i-1}^2 u_i^2$, $v_{i-1}^3 u_i^3$, for $i \geq 2$.





Figure 3: Segment graph G_T

Let $T_1(n)$ be the graph obtained from $T(n)$ by adding the edges $u_1^1 v_n^1$, $u_1^2 v_n^2$, $u_1^3 v_n^3$. Let $T_2(n)$ be the graph obtained from $T(n)$ by adding the edges $u_1^3 v_n^1$, $u_1^2 v_n^2$, $u_1^1 v_n^3$. Let $T_3(n)$ be the graph obtained from $T(n)$ by adding the edges $u_1^1 v_n^3$, $u_1^2 v_n^1$, $u_1^3 v_n^2$. In [4], Boben proved that for each fixed value of n , no two of the graphs $T_1(n)$, $T_2(n)$, $T_3(n)$ are isomorphic.

Note that $T_1(1)$ is the Levi graph of Desargues' configuration, and $T_2(1)$, $T_3(1)$ correspond to the Levi graphs of the configurations 10_3F and 10_3G respectively according to Kantor's [9] notation for the ten 10_3 configurations.

The Pappus graph: Recall that the Levi graph of the Pappus 9_3 configuration is the following pseudo 2-factor isomorphic but not 2-factor isomorphic cubic bipartite graph [2], called the *Pappus graph* P_0 .



3 2-factors of Irreducible Levi Graphs

Let G be a graph and u, v be two vertices in G . Then a (u, v) -path is a path from u to v . Given two disjoint paths $P = u_1, \dots, u_n$ and $Q = u_{n+1}, \dots, u_{n+m}$ (except maybe for $u_1 = u_{n+m}$), the path $PQ = u_1, \dots, u_{n+m}$ is the concatenation of P and Q together with the edge $u_n u_{n+1}$. Similarly, for a vertex $v \in (G - P) \cup \{v_1\}$, the path Pv is composed by P , v and the edge $u_n v$. If $u_1 = u_{n+m}$ or $u_1 = v$ we write (PQ) and (Pv) respectively, to emphasize that PQ and Pv are circuits.

Theorem 3.1 *The Heawood and the Pappus graphs are the only irreducible Levi graphs which are pseudo 2-factor isomorphic.*

Proof. It is straightforward to show that the Heawood graph H_0 is 2-factor hamiltonian and hence pseudo 2-factor isomorphic (cf. [7]). We have already proved in [2, Proposition 3.3] that the Pappus graph is pseudo 2-factor isomorphic. We need to prove that all other irreducible Levi graphs are not pseudo 2-factor isomorphic and we will do so by finding two 2-factors with different parity of number of circuits in each of them.

The following paths will be used for constructing 2-factors in $D(n)$, for $n \geq 8$.

$$\begin{aligned} L_i^1 &= u_i^1 u_i^2 u_i^3 u_i^4 v_i^2 v_i^1 & L_i^2 &= u_i^4 u_i^3 u_i^2 u_i^1 v_i^1 v_i^2 \\ M_1 &= u_1^4 v_1^2 u_2^4 u_3^2 u_3^3 u_3^4 v_2^2 v_2^1 u_3^1 v_3^1 v_3^2 & M_2 &= u_m^4 v_m^2 v_m^1 u_m^1 u_m^2 u_m^3 w_m^1 w_m^2 u_1^4 \\ N_i &= u_i^2 u_i^1 v_{i-1}^1 v_{i-1}^2 u_i^4 u_i^3 & N_m &= w_m^2 w_m^1 v_m^1 v_m^2 w_m^4 w_m^3 u_1^4 v_1^2 u_2^4 u_3^3 \\ C_1 &= (u_1^1 u_1^2 u_1^3 u_2^2 u_2^1 u_1^1) & C_2 &= (u_1^1 u_1^2 u_1^3 u_1^4 w_m^2 w_m^1 u_1^1) \\ C_3 &= (v_1^1 v_1^2 u_2^4 v_2^2 v_2^1 u_3^1 v_3^1 v_3^2 u_3^4 u_3^3 u_3^2 u_2^2 u_2^1 v_1^1) \end{aligned}$$


$$\text{Hamiltonian 2-factors in } D(n) \text{ are } \begin{cases} (L_1^1 L_2^1 \cdots L_m^1 u_1^1) & n \equiv 0 \pmod{3} \\ (L_1^1 L_2^1 \cdots L_m^1 w_m^2 w_m^1 u_1^1) & n \equiv 1 \pmod{3} \\ (L_1^1 L_2^1 \cdots L_m^1 w_m^1 w_m^2 w_m^3 w_m^4 u_1^1) & n \equiv 2 \pmod{3} \end{cases}$$

Disconnected 2-factors with exactly two circuits in $D(n)$ are

$$\left\{ \begin{array}{lll} C_1 \cup (M_1 u_1^4) & n = 9 & \text{for } n \equiv 0 \pmod{3} \\ C_1 \cup (M_1 L_4^2 \cdots L_m^2 u_1^4) & n = 3m, m \geq 4 & \\ \\ C_2 \cup C_3 & n = 10 & \\ C_1 \cup (M_1 M_2) & n = 13 & \text{for } n \equiv 1 \pmod{3} \\ C_1 \cup (M_1 L_4^2 \cdots L_{m-1}^2 M_2) & n = 3m + 1, m \geq 5 & \\ \\ C_1 \cup (N_m w_m^2) & n = 8 & \text{for } n \equiv 2 \pmod{3} \\ C_1 \cup (N_3 \cdots N_m u_3^2) & n = 3m + 2, m \geq 3 & \end{array} \right.$$

Now we need to find such pairs of 2-factors for the graphs $T_1(n)$, $T_2(n)$ and $T_1(n)$, $n \geq 1$. To this purpose we need to consider the following six paths in the segment graph G_T from Fig 3.

$$\begin{aligned}
P_i^1 &= u_i^2, w_i^1, u_i^1, w_i^2, u_i^3, w_i^3, x_i^3, y_i^3, z_i^3, v_i^3, t_i^2, v_i^1, z_i^1, y_i^2, x_i^2, t_i^1, x_i^1, y_i^1, z_i^2, v_i^2 \\
P_i^2 &= u_i^2, w_i^1, u_i^1, w_i^2, u_i^3, w_i^3, x_i^3, y_i^3, z_i^2, v_i^2 \\
(P_i^3) &= (v_i^1, t_i^2, v_i^3, z_i^3, y_i^2, x_i^2, t_i^1, x_i^1, y_i^1, z_i^1, v_i^1) \\
Q_i^1 &= u_i^3, w_i^3, u_i^2, w_i^1, u_i^1, w_i^2, x_i^2, y_i^2, z_i^3, v_i^3, t_i^2, v_i^1, z_i^1, y_i^1, x_i^1, t_i^1, x_i^3, y_i^3, z_i^2, v_i^2 \\
Q_i^2 &= u_i^3, w_i^2, u_i^1, w_i^1, u_i^2, w_i^3, x_i^3, y_i^3, z_i^2, v_i^2
\end{aligned}$$



The paths P_i^1 and Q_i^1 are hamiltonian (u_i^2, v_i^2) and (u_i^3, v_i^2) -paths, respectively. The paths P_i^2 and Q_i^2 are (u_i^2, v_i^2) , and (u_i^3, v_i^2) -paths on 10 vertices, respectively. Finally, (P_i^3) is a 10-circuit in $G_T - P_i^2 = G_T - Q_i^2$.

In $T_1(n)$ and $T_2(n)$ the hamiltonian 2-factor $F_1(n) = (P_1^1 P_2^1 \cdots P_n^1 u_1^2)$ and the disconnected 2-factor with exactly two circuits $F_2(n) = (P_1^2 P_2^1 \cdots P_n^1 u_1^2) \cup (P_1^3)$ (even) show that these graphs are not pseudo 2-factor isomorphic.

Similarly, in $T_3(n)$ the hamiltonian 2-factor $F'_1(1) = (Q_1^1 P_2^1 \cdots P_n^1 u_1^3)$ and the disconnected 2-factor with exactly two circuits $F'_2(1) = (Q_1^2 P_2^1 \cdots P_n^1 u_1^3) \cup (P_1^3)$ show that these graphs are not pseudo 2-factor isomorphic. $\square$

Note that Theorem 3.1 proves Conjecture 1.1 in the case of irreducible pseudo 2-factor isomorphic cubic bipartite graphs. In the next Section we show that Conjecture 1.1 cannot be directly proved from Theorem 3.1 by extending it to reducible Levi graphs.

4 2-factors in extensions and reductions of Levi graphs of n_3 configurations

Recall that a *Martinetti extension* can be described in terms of graphs as follows:

Let G_1 be the Levi graph of a symmetric configuration n_3 and suppose that in G_1 there are two edges $e_1 = x_1 y_1$ and $e_2 = x_2 y_2$ with no common neighbours, then

the graph $G := G_1 - \{e_1, e_2\} + \{u, v\} + \{ux_1, ux_2, vy_1, vy_2\}$ is the Levi graph of an $(n+1)_3$ configuration.




Figure 4: Martinetti Extension

Similarly the Levi graph G of a symmetric configuration $(n+1)_3$ is *Martinetti reducible* if there is an edge $e = uv$ in G such that either $G := G_1 - \{u, v\} + x_1y_1 + x_2y_2$ or $G := G_1 - \{u, v\} + x_1y_2 + x_2y_1$ is again the Levi graph of a symmetric configuration n_3 , where x_1, x_2, y_1, y_2 are the neighbours of u and v as in the following figure:




Figure 5: Martinetti Reduction

It is well known that the 7_3 configuration, whose Levi graph is the Heawood graph, is not Martinetti extendible and that the Pappus configuration is Martinetti extendible in a unique way; it is easy to show that this extension is not pseudo 2-factor isomorphic.

Let $\mathcal{C}$ be a symmetric configuration n_3 and $\mathcal{C}'$ be a symmetric configuration $(n+1)_3$ obtained from $\mathcal{C}$ through a Martinetti extension. It can be easily checked that there are 2-factors in $\mathcal{C}'$ that cannot be reduced to a 2-factor in $\mathcal{C}$. For example if $\mathcal{C}$ corresponds to the first option in Fig. 5, a 2-factor of $\mathcal{C}'$ containing the path x_1uvy_2 will not reduce to a 2-factor in $\mathcal{C}$. Conversely, there might be 2-factors of $\mathcal{C}$ for which the parity of number of circuits is not preserved when extended to a 2-factor in $\mathcal{C}'$. For example, the graph $H_0 * H_0$ (the star product [8, p. 90] of the Heawood graph with itself) which is 2-factor hamiltonian and Martinetti reducible (only through the edges of the non-trivial 3-edge-cut), has all Martinetti reductions which are no longer pseudo 2-factor isomorphic. Hence, we cannot directly prove Conjecture 1.1 by studying the 2-factors of reducible configurations from the set of 2-factors of their underlying irreducible ones.

References

- [1] M. Abreu, D.Labbate, J. Sheehan. Pseudo and Strongly Pseudo 2-factor isomorphic Regular Graphs, (submitted).

- [2] M. Abreu, A. Diwan, B. Jackson, D. Labbate, J. Sheehan. Pseudo 2-factor isomorphic Regular Bipartite Graphs, *J. Combin. Theory Ser. B* **98** (2008) 432–442.
- [3] R. Aldred, M. Funk, B. Jackson, D. Labbate and J. Sheehan, Regular bipartite graphs with all 2-factors isomorphic, *J. Combin. Th. Ser. B*, 92 (2004), no. 1, 151–161.
- [4] M. Boben, Irreducible (v_3) configurations and graphs, *Discrete Mathematics*, **307** (2007) 331–344.
- [5] A.A. Diwan, Disconnected 2-factors in planar cubic bridgeless graphs, *J. Combin. Th. Ser. B*, 84, (2002), 249–259.
- [6] R.J. Faudree, R.J. Gould, and M.S. Jacobson, On the extremal number of edges in 2-factor hamiltonian graphs, *Graph Theory - Trends in Mathematics*, Birkhäuser (2006), 139–148.
- [7] M. Funk, B. Jackson, D. Labbate and J. Sheehan, 2-factor hamiltonian graphs, *J. of Combin. Th. Ser. B*, 87, (2003), no.1, 138–144.
- [8] D.A. Holton and J. Sheehan, *The Petersen graph*, Australian Mathematical Society Lecture Series, 7. Cambridge University Press, Cambridge, 1993.
- [9] S. Kantor, Die Configurationen $(3, 3)_{10}$, *Sitzungsber. Wiener Akad.* **84** (1881), 1291–1314.
- [10] V. Martinetti, Sulle configurazioni piane μ_3 , *Annali di Matematica Pura ed Applicata*, Ser. II - 15 (dall’aprile 1867 al gennaio 1888) 1–26.